

# YOU'RE TAKING THE NEXT STEP WITH NUBEQA®

Pronounced *New-Becca*

  
**NUBEQA**  
(darolutamide) 300 mg tablets

## HOW DOES NUBEQA WORK?

NUBEQA helps **delay the growth of prostate cancer cells** by blocking testosterone from attaching to a molecule called an **androgen receptor**.

## HOW SHOULD I TAKE NUBEQA?

- Take NUBEQA exactly as your doctor tells you
- Your doctor may change your dose if needed. Do not stop taking your prescribed dose of NUBEQA without talking with your doctor first
- Swallow two tablets whole, 2 times a day, with food
- Continue on hormone therapy while taking NUBEQA
- If you take more NUBEQA than prescribed, call your doctor right away



## WHAT SHOULD I DO IF I MISS A DOSE OF NUBEQA?

- Take your prescribed dose as soon as you remember
- Do not take 2 doses together to make up for a missed dose

## WHAT IS NUBEQA®?

NUBEQA® (darolutamide) is a prescription medicine used to treat adults with prostate cancer:

- that has not spread to other parts of the body and no longer responds to medical or surgical treatment that lowers testosterone (non-metastatic castration-resistant prostate cancer, or nmCRPC), **or**
- that has spread to other parts of the body and responds to medical or surgical treatment that lowers testosterone (metastatic hormone-sensitive prostate cancer, or mHSPC), in combination with docetaxel

It is not known if NUBEQA is safe and effective in women or children.

## IMPORTANT SAFETY INFORMATION

**Before taking NUBEQA, tell your healthcare provider about all your medical conditions, including if you:**

- have heart problems, high blood pressure, diabetes, or high amounts of fat or cholesterol in your blood (dyslipidemia)
- have a history of seizures, brain injury, stroke, or brain tumors
- have kidney or liver problems
- are pregnant or plan to become pregnant. NUBEQA can cause harm to your unborn baby and loss of pregnancy (miscarriage)
- have a partner who may become pregnant. Males who have female partners who may become pregnant should use effective birth control (contraception) during treatment and for 1 week after the last dose of NUBEQA. Talk with your healthcare provider about birth control methods

Please see additional Important Safety Information on the back page and the full [Prescribing Information](#).

## IMPORTANT SAFETY INFORMATION (CONTINUED)

**Before taking NUBEQA, tell your healthcare provider about all your medical conditions, including if you (continued):**

- are breastfeeding or plan to breastfeed. It is not known if NUBEQA passes into breast milk

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. NUBEQA may affect the way other medicines work and other medicines may affect how NUBEQA works.

You should not start or stop any medicine before you talk with the healthcare provider that prescribed NUBEQA. Know the medicines you take. Keep a list of them with you to show to your healthcare provider and pharmacist when you get a new medicine.

### **NUBEQA may cause serious side effects, including:**

- **Heart disease.** Blockage of the arteries in the heart (ischemic heart disease) that can lead to death has happened in some people during treatment with NUBEQA. Your healthcare provider will monitor you for signs and symptoms of heart problems during your treatment with NUBEQA. Call your healthcare provider or get medical help right away if you get chest pain or discomfort at rest or with activity, or shortness of breath during your treatment with NUBEQA
- **Seizure.** Treatment with NUBEQA may increase your risk of having a seizure. You should avoid activities where a sudden loss of consciousness could cause serious harm to yourself or others. Tell your healthcare provider right away if you have loss of consciousness or seizure

### **The most common side effects of NUBEQA in people with nmCRPC include:**

- increase in liver function tests
- decreased white blood cells (neutropenia)
- feeling more tired than usual
- arm, leg, hand, or foot pain
- rash

### **The most common side effects of NUBEQA when used in combination with docetaxel in people with mHSPC include:**

- constipation
- rash
- decreased appetite
- bleeding
- weight gain
- high blood pressure
- decreased red blood cells (anemia)
- high blood sugar levels
- decreased white blood cells
- increase in liver function tests
- low blood calcium levels

NUBEQA may cause fertility problems in males, which may affect the ability to father children. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of NUBEQA.

Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

### **CALL TO TALK WITH A SAVINGS AND SUPPORT SPECIALIST TODAY**

**1-800-288-8374**

Monday-Friday, 8:00 AM - 8:00 PM (ET)

 **Access Services**  
by Bayer™

### **HEAR MORE FROM A REAL NUBEQA PATIENT**



**WATCH NOW**

Please see additional Important Safety Information on the front page and the full [Prescribing Information](#).



© 2025 Bayer. All rights reserved. BAYER, the Bayer Cross, and NUBEQA are registered trademarks of Bayer, and Access Services by Bayer is a trademark of Bayer. PP-NUB-US-3853-1 01/2025

  
**NUBEQA**<sup>®</sup>  
(darolutamide) 300 mg  
tablets